

A study of the tunnelling-charging Hamiltonian of a Cooper pair pump

M. Aunola

Dept. of Physics, University of Jyväskylä, P.O. Box 35 (Y5), FIN-40351 Jyväskylä, Finland

Email: Matias.Aunola@phys.jyu.fi

(February 6, 2008)

General properties of the tunnelling-charging Hamiltonian of a Cooper pair pump are examined with emphasis on the symmetries of the model. An efficient block-diagonalisation scheme and a compatible Fourier expansion of the eigenstates is constructed and applied in order to gather information on important observables. Systematics of the adiabatic pumping with respect to all of the model parameters are obtained and the link to the geometrical Berry's phase is identified.

Adiabatic transport of single electrons in arrays of small metallic tunnel junctions has been widely studied in recent years.^{1,2} In the Coulomb blockade regime phase-shifted gate voltages have been used to induce a dc current $I = -nef$, where n is the number of carried electrons and f is the gating frequency. Normal-state pumps transporting single electrons have reached accuracy that can be considered for metrological applications.² Pumping of Cooper pairs has gained interest due to new ideas in quantum measuring and computing.^{3,4}

A quantitative theory of pumping Cooper pairs in gated arrays of Josephson junctions when the environment has negligible impedance has been presented.^{5,6} The leading order pumped current is $I \approx -2ef[1 - a_N(\varepsilon_J)^{N-2} \cos \phi]$, where a_N is a constant and $\varepsilon_J := E_J/E_C$ is the coupling strength. Here E_J and $E_C := (2e)^2/2C$ are the Josephson coupling energy and the charging energy, respectively. According to recent calculations the $\cos \phi$ -dependent inaccuracy should be experimentally observable at least in a certain frequency range.⁷ In this article the model for adiabatic pumping from Refs. 5 and 6 is examined in a thorough manner, concentrating on the symmetries and general features of the simplified system, thus extending the treatment to strong couplings and long arrays. The overall behaviour of the pumped current is explained and systematised, but not rigorously proven.

A schematic view of a Cooper pair pump and the ideal operation of the gate voltages $V_{g,k}$ are shown in Fig. 1. On any of the N legs of a cycle at most two gate voltages are changed. The tunnelling-charging Hamiltonian

$$H = H_C + H_J, \quad (1)$$

neglects the quasiparticle tunnelling and other degrees of freedom. The full set of model parameters are ε_J , total phase difference over the array ϕ , the relative junction capacitances $\vec{c}$, where $c_k := C_k/C$ and $\sum_{k=1}^N C_k^{-1} = N/C$, and the (normalised) gate charges $\vec{q} := \{q_1, \dots, q_{N-1}\}$, where $q_k := -C_{g,k}V_{g,k}/2e$. For homogeneous arrays $c_k := 1$ and the inhomogeneity is quantified by $X_{\text{inh}} =$

$[\sum_k (c_k^{-1} - 1)^2/N]^{1/2}$. The model Hamiltonian is diagonal with respect to ϕ , which is assumed to be fixed, but could be controlled by an external bias voltage according to $d\phi/dt = -2eV/\hbar$ and is subject to any voltage fluctuations.^{7,8} The conjugate variable $\hat{M}$, the average number of tunnelled Cooper pairs, is undetermined in the present model.




FIG. 1. (a) A schematic drawing of a gated Josephson array of $N (\geq 3)$ junctions. The C_k and $C_{g,k}$ are the capacitances of the junctions and gates, respectively. b) The optimal operation of the gate charges $\vec{q}$. Each cycle carries approximately one Cooper pair through the array, when $\varepsilon_J \ll 1$.

The matrix elements of the charging Hamiltonian H_C are given by the capacitive charging energy

$$\langle \vec{n} | H_C(\vec{q}) | \vec{n} \rangle_\phi = E_C \left[\sum_{k=1}^N \frac{v_k^2}{c_k} - \frac{1}{N} \left(\sum_{k=1}^N \frac{v_k}{c_k} \right)^2 \right], \quad (2)$$

where the number of Cooper pairs on each island is given by $\vec{n}$. The quantities v_k , $k = 1, \dots, N$, are a solution of

$$v_k - v_{k+1} = n_k - q_k. \quad (3)$$

Tunnelling of one Cooper pair through the k th junction changes $|\vec{n}\rangle$ by $\vec{\delta}_k$, where the non-zero components are (if applicable) $(\vec{\delta}_k)_k = 1$ and $(\vec{\delta}_k)_{k-1} = -1$. The tunnelling Hamiltonian then reads

$$H_J = - \sum_{\vec{n}, k=1}^N \frac{c_k E_J}{2} (|\vec{n} + \vec{\delta}_k\rangle \langle \vec{n}| e^{i\phi/N} + \text{H.c.}). \quad (4)$$

The supercurrent flowing through the array is determined by the supercurrent operator

$$I_S = (-2e/\hbar)(\partial H/\partial \phi), \quad (5)$$

a Gâteaux derivative⁹ of the full Hamiltonian. By changing the gate voltages adiabatically along a closed path Γ , a charge transfer $Q_{\text{tot}} := Q_s + Q_p$ is induced. The pumped charge, Q_p , depends only on the chosen path, while the charge due to direct supercurrent, Q_s , also depends on how the gate voltages are operated. If the system remains in a adiabatically evolving state $|m\rangle$, the total transferred charge, Q_{tot} , in units of $-2e$ reads^{5,6}

$$\frac{1}{\hbar} \int_0^\tau \frac{\partial E_m(t)}{\partial \phi} dt - 2 \oint_\Gamma \sum_{l \neq m} \text{Im} \left[\frac{\langle m | I_S | l \rangle \langle l | dm \rangle}{E_l - E_m} \right], \quad (6)$$

where $|dm\rangle$ is the change in $|m\rangle$ due to a differential change of the gate voltages $d\vec{q}$.

Intermediate states $|l\rangle$ and energy denominators can be removed by rewriting the off-diagonal matrix element between stationary states as

$$\langle m | \partial H / \partial \phi | l \rangle = (E_l - E_m) [\partial(\langle m |) / \partial \phi] | l \rangle, \quad (7)$$

and using the identity $\sum_l |l\rangle \langle l| = \hat{1}$. The canonical representation $\hat{M} = -i\partial/\partial\phi$ connects the pumped charge to the average number of tunnelled Cooper pairs by

$$Q_p = 2 \oint_\Gamma \text{Re} \left[\langle m | \hat{M} | dm \rangle \right]. \quad (8)$$

A single eigenstate for phase differences ϕ and $\phi + d\phi$ is required for each integration point. This expression identifies $\hat{M}$ as the link between Q_p and the geometrical Berry's phase, $\gamma_m(\Gamma) = i \oint_\Gamma \langle m | dm \rangle$.¹⁰ The connection was given without an explicit identification in Ref. 5 and was mentioned in Ref. 11. For $\varepsilon_J \rightarrow 0$ only two charge are of importance during each leg. From the wave function $|m\rangle = [(1-a^2)^{1/2}, ae^{i\phi/N}]^T$, where $a: 0 \rightarrow 1$, one obtains $\langle m | \hat{M} | dm \rangle = d(a^2)/2N$ and $Q_p = 1$ for a full cycle.

A reference state $|\vec{n}_0\rangle$ induces a convenient labelling of the charge states. Each state is denoted by integers $\{y_k\}_{k=1}^N$, $0 \leq Y_{\vec{n}} := \sum_k y_k < N$, such that $\vec{n} = \vec{n}_0 + \sum_k y_k \vec{\delta}_k$. The numbers y_k tabulate the number and direction of tunnellings from $\vec{n}_0$ to $\vec{n}$. The distance between charge states is defined by $\tilde{d}(\vec{n}_1, \vec{n}_2) := \min \left(\sum_{k=1}^N |y_k^{(1)} - y_k^{(2)} + l| : l \in \mathbb{Z} \right)$. If $\tilde{d}(\vec{n}_1, \vec{n}_2) = 1$ ($=l$), then $|\vec{n}_1\rangle$ and $|\vec{n}_2\rangle$ are (l th) nearest neighbours.

A change of basis $U\{|\vec{n}\rangle\} = \{|e^{i\phi Y_{\vec{n}}/N} \vec{n}\rangle\}$ yields a Hamiltonian matrix $\tilde{H} = U H U^{-1}$ with properties $\tilde{H}(\phi + 2\pi) = \tilde{H}(\phi)$ and $\tilde{H}(-\phi) = (\tilde{H}(\phi))^*$. Thus eigenstates and eigenvalues of $\tilde{H}$ are periodic under 2π , or state labels change cyclically. Usually the former happens, so the supercurrent in a stationary state is described by a Fourier sine series

$$\langle I_S \rangle_{(N, \varepsilon_J, \phi, \vec{c}, \vec{q})} := \sum_{l=1}^\infty \alpha_l \sin(l\phi). \quad (9)$$

The ground state supercurrent behaves differently only at the so-called resonance points, where the ground state

becomes degenerate for $\phi = \pi + 2l\pi$, $l \in \mathbb{Z}$. For homogeneous arrays resonance points are located at $\vec{q} = \vec{n} \pm (1, \dots, 1)/N$, where $\vec{n}$ is arbitrary. The corresponding poles in the Berry's phase give raise to Q_p , and ε_J determines which poles are important.

Due to symmetry of the representation $\tilde{H}$ the original amplitudes in 2π -periodic states are given by

$$a_{\vec{n}}^{\vec{q}, \phi} = \sum_{l=-\infty}^\infty a_{\vec{n}, l}^{\vec{q}} e^{i\phi(l + Y_{\vec{n}}/N)}, \quad (10)$$

where real Fourier coefficients $a_{\vec{n}, l}^{\vec{q}}$ are fixed by the gauge condition $a_{\vec{n}'}^{\vec{q}, \phi} := |a_{\vec{n}'}^{\vec{q}, \phi}| e^{i\phi Y_{\vec{n}'}/N}$ for the charge state $|\vec{n}'\rangle$. The averaged number of tunnelled Cooper pairs,

$$\mathcal{M} := \sum_{\vec{n}, l} (l + Y_{\vec{n}}/N) (a_{\vec{n}, l}^{\vec{q}})^2, \quad (11)$$

is unique up to a gauge-dependent integer. The apparent contradiction between sharp phase difference combined with sharp value of $\mathcal{M}$ is an artefact due to gauge-fixing. Stronger coupling increases the variance of $\mathcal{M}$. Discontinuities in gauges certainly occur on any closed path encircling an odd number of resonance points. A gauge is unstable near a discontinuity, but everywhere away from resonance points many valid gauges exist. Especially on the gating path depicted in Fig. 1, the dominant charge states on each leg give very stable gauges.

The pumped charge, Q_p , can be evaluated using a gauge-independent differential expression

$$dQ_p(\phi) = \sum_{l'=0}^\infty \sum_{\vec{n}, l=-\infty}^\infty \left[\frac{2(l + Y_{\vec{n}}/N)}{1 + \delta_{l'0}} d(a_{\vec{n}, l} a_{\vec{n}, l+l'}) + l' (a_{\vec{n}, l} da_{\vec{n}, l+l'} - a_{\vec{n}, l+l'} da_{\vec{n}, l}) \right] \cos(l'\phi), \quad (12)$$

where the ϕ -independent average is simply $d\mathcal{M}$. Due to the normalisation of $|m\rangle_\phi$, the coefficients are orthonormal, i.e. $\sum_{\vec{n}, l} a_{\vec{n}, l} a_{\vec{n}, l+l'} = \delta_{l'0}$, which cancels the terms multiplying the full differential by l' . Expression (12) indicates that the averaged charge transfer for a full cycle is exactly $-2e$, regardless of the inhomogeneity of the array or reasonable deformations of the gating path.

The tunnelling-charging Hamiltonian can often be block diagonalised with the following transformation. Let an orthonormal basis $\{|s\rangle\}$ span the Hilbert space $\mathcal{H}$ and the matrix elements of a Hamiltonian H be $h_{ss'} := \langle s | H | s' \rangle$. Choose projection operators $\{P_i\}$ by $P_i = \sum_{k=1}^{d_i(<\infty)} |i_k\rangle \langle i_k|$ and require that $\sum_{ij} P_i P_j = \sum_i P_i = \hat{1}$. If all of the row sums

$$W_{ij, k} := \sum_{k'=1}^{d_j} h_{i_k j_{k'}}, \quad k = 1, \dots, d_i, \quad (13)$$

are independent of k , the Hamiltonian commutes with the projection operator $P := \sum_i |\psi_i\rangle \langle \psi_i|$, where $|\psi_i\rangle = d_i^{-1/2} \sum_{k=1}^{d_i} |i_k\rangle$. Thus H can be written as a direct sum $H = H_P \oplus H_{\hat{1}-P}$ with matrix elements of H_P given as

$$h_{ij} := \langle \psi_i | H | \psi_j \rangle = W_{ij} (d_i/d_j)^{1/2} = W_{ji}^* (d_j/d_i)^{1/2}. \quad (14)$$

The block-diagonalising transformation is not specific to the present problem, as it amounts to extracting states with a specific symmetry from all of the system's eigenstates. The ground state of $H_C + H_J$ is always an eigenstate of H_P , because $a_{\vec{n}}^{\vec{q}, \phi=0} > 0$ for all $\vec{q}$ and $\vec{n}$. The Fourier expansion (10) can be used simultaneously if each subspace label, i , corresponds to a single value of $Y_{\vec{n}}$.

For homogeneous arrays at $\vec{q} = \vec{n}_0$ all junctions are indistinguishable in terms of the charging energy. The subspace labels are of the form $\bar{m}_{\bar{z}}$ with $d_{\bar{m}_{\bar{z}}} = N! / (\prod_{j=1}^{j_{\max}} m_j!)$. This subspace contains charge eigenstates carrying the label

$$\vec{y} = \left(z_1^{(1)}, \dots, z_1^{(m_1)}, z_2^{(1)}, \dots, z_2^{(m_2)}, \dots, z_{j_{\max}}^{(m_{j_{\max}})} \right), \quad (15)$$

where $z_1 < \dots < z_{j_{\max}}$ and $m_1 + \dots + m_{j_{\max}} = N$, or any distinct permutation of $\vec{y}$. Thus only the number and multiplicity of tunnellings is of importance.

On the gating path the gate charges can be written as $\vec{q} = (1-x)\vec{n}_0 + x\vec{n}_1$, where $|\vec{n}_0\rangle$ ($|\vec{n}_1\rangle$) is the initial (final) optimal charge state for the given leg. Thus one junction becomes distinguishable and subspace labels can be chosen as $z_0; \bar{m}_{\bar{z}}$, where $\bar{m}_{\bar{z}}$ refers to the remaining $N-1$ components, $0 \leq z_0 + \sum_{i=1}^{N-1} m_i z_i < N$ and $d_{z_0; \bar{m}_{\bar{z}}} = (N-1)! / (\prod_{j=1}^{j_{\max}} m_j!)$.

The diagonal and non-zero off-diagonal matrix elements of H_P are the common charging energies and

$$- (\varepsilon_J m_j / 2) e^{\pm i \phi / N} (d_{(z_0; \bar{m}_{\bar{z}})} / d_{(z'_0; \bar{m}'_{\bar{z}})})^{1/2} \quad (16)$$

with $m_0 = 1$ when applicable, respectively. The block-diagonalised matrices are sparser than the original matrices and the amplitudes in eigenvectors are multiplied by $(d_{(z_0; \bar{m}_{\bar{z}})})^{1/2}$ due to combining of several amplitudes into one. A further block diagonalisation is possible at $\vec{q} = \vec{n}_0$ and $\vec{q} = (\vec{n}_0 + \vec{n}_1)/2$ if ϕ is a multiple of π .

A truncation picks the charge states required for reliable evaluation of eigenstates and observables. First an initial truncation, a set of states $B_i = \{|\vec{n}_j\rangle\}$, is chosen. Its extensions, bases $B_i^{(l)}$, contain all neighbours up to and including l th nearest neighbours of each and every $|\vec{n}_j\rangle$. For non-ideal cycles and/or inhomogeneous arrays the initial truncation is the “b basis” of Ref. 6 which reproduces leading order supercurrent and inaccuracy. A “c basis” is just the first order extension of a “b basis”, i.e. “b basis⁽¹⁾”.

For ideal cycles and homogeneous arrays the optimal B_0 truncation corresponds to $3N-2$ labels, that is $0; (j, N-1-j)_{(0,1)}$, “1; $(j, N-1-j)_{(-1,0)}$ ”, $j = 0, \dots, N-1$, and $1; (j, N-1-j)_{(0,1)}$, $j = 1, \dots, N-2$. These $3 \cdot 2^{N-1} - 2$ charge states actually contribute to the leading order supercurrent and inaccuracy. More restricted, few-state truncations $B_1 = \{|\vec{n}_0\rangle\}$ and $B_2 = \{|\vec{n}_0\rangle, |\vec{n}_1\rangle\}$ are of use especially when the ground state energy is sought. The efficiency of the block diagonalisation for bases $B_0^{(l)}$ is shown in Table I where the number of labels is compared against the number of included charge eigenstates. Examples of bases $B_1^{(l)}$ and $B_2^{(l)}$ are also given.

TABLE I. The number of subspace labels (sl) and included charge states (ch) for bases $B_0^{(l)}$ with several N and examples of bases $B_1^{(l)}$ and $B_2^{(l)}$.

$N_{0/1/2}$	sl _{<i>l</i>}	ch	sl _{<i>l</i>}	ch
4 ₀	52 ₂	182	1872 ₁₄	9572
5 ₀	82 ₂	626	3082 ₁₁	46400
6 ₀	114 ₂	1918	3468 ₉	$1.56 \cdot 10^5$
7 ₀	146 ₂	5428	3941 ₈	$5.22 \cdot 10^5$
8 ₀	178 ₂	14498	3648 ₇	$1.30 \cdot 10^6$
9 ₀	210 ₂	37082	2899 ₆	$2.44 \cdot 10^6$
8 ₁	69 ₅	12331	14727 ₁₉	$6.15 \cdot 10^7$
7 ₂	248 ₅	9452	12842 ₁₄	$2.19 \cdot 10^6$

As a concrete example, take basis $B_1^{(l)}$ and the subspace labels N_0 , $(N-1, 1)_{(0,1)}$ and $(1, N-1)_{(0,1)}$, standing for $2N+1$ charge eigenstates at $\vec{q} = \vec{n}_0$. The block-diagonalised Hamiltonian obtained from Eqs. (14) and (16) is given by

$$H_{\text{bd}} = \begin{pmatrix} 0 & K_N^* & K_N \\ K_N & (N-1)/N & 2\delta_{N3} K_N^* / \sqrt{N} \\ K_N^* & 2\delta_{N3} K_N / \sqrt{N} & (N-1)/N \end{pmatrix} \quad (17)$$

where $K_N := -(\varepsilon_J/2)e^{i\phi/N}\sqrt{N}$. For $N=3$ the leading order supercurrent for $\varepsilon_J \ll 1$ is already reproduced. Note that labels $(N-1, 1)_{(0,1)}$ and $(1, N-1)_{(0,1)}$ can be joined, if ϕ is a multiple of π .

The discussion has now lead to the main results of the paper, the general systematics of the pumped charge. For ideal gating sequences $d\mathcal{M} = 1/N$ for each leg due to symmetry. Corresponding integrated pumped charge reads

$$Q_p(N, \varepsilon_J, \phi, \vec{c}, \text{leg}) := 1/N + \sum_{l=1}^{\infty} b_{l, \text{leg}} \cos(l\phi), \quad (18)$$

which is always positive. Replace $1/N$ by $d\mathcal{M}$ for non-ideal cycles. When performing numerical calculations, the basis must contain all of the important charge states and the number of angles used in fast Fourier transform, 2^j , must be large enough not to allow misidentification of non-negligible components $a_{\vec{n}, l}$ and $a_{\vec{n}, l+2j}$ as a single component. Failure to satisfy these requirements causes systematical error due to incorrect amplitudes and spurious interference between Fourier components in Eq. (10), respectively. The integration points should be chosen so that the magnitude of norms $\|dm\|$ is neither too large nor varies too much.

The rate of convergence for each $\cos(l'\phi)$ -dependent term in Eq. (18) behaves as $1/(\# \text{steps})^2$. For a reasonable choice of integration points 300–500 steps per leg, much less than used in Refs. 5 and 6, usually gives relative precision of the order of 10^{-5} (disregarding the systematical error) for coefficients with magnitude greater than 10^{-8} . This greatly diminishes convergence problems due

to smallness of $da_{\vec{n},l}$ for large bases and large number of angles. The expression $2l'a_{\vec{n},l}da_{\vec{n},l+l'}$ has been used as the latter part of Eq. (12) for many of the data points. In this case the rate of convergence behaves as $1/\#\text{steps}$. Bases $B_2^{(l)}$ have also been used when calculating Q_p .

TABLE II. The first four ratios β_l and the limiting value $\tilde{\beta}$ for $\varepsilon_J = 0.4$ as function of N and for $N = 4$ as function of ε_J . Subscript of $\tilde{\beta}$ in the last column indicates the coefficient β_l used when estimating the limiting value.

$(\varepsilon_J)_N$	β_1	β_2	β_3	β_4	$(\tilde{\beta})_l$
0.4 ₃	0.8802	0.7782	0.7470	0.7347	0.722 ₆₀
0.4 ₄	0.7250	0.5706	0.5354	0.5225	0.509 ₂₅
0.4 ₅	0.5494	0.3842	0.3551	0.3451	0.335 ₂₀
0.4 ₆	0.3844	0.2393	0.2188	0.2121	0.206 ₁₅
0.4 ₇	0.2516	0.1401	0.1272	0.1231	0.120 ₁₁
0.4 ₈	0.1563	0.0785	0.0708	0.0685	0.067 ₉
0.4 ₉	0.0934	0.0427	0.0383	0.0370	0.036 ₇
0.4 ₁₀	0.0542	0.0227	0.0203	0.0196	0.019 ₇
0.04 ₄	0.0188	0.0096	0.0095	0.0095	0.0095 ₄
0.2 ₄	0.3444	0.2164	0.2049	0.2025	0.201 ₁₄
0.6 ₄	0.8853	0.7833	0.7469	0.7296	0.701 ₄₀
0.8 ₄	0.9475	0.8876	0.8598	0.8444	0.808 ₈₀
1.0 ₄	0.9738	0.9390	0.9198	0.9079	0.870 ₁₂₀
1.2 ₄	0.9860	0.9654	0.9526	0.9441	0.908 ₁₆₀

The sequence of the Fourier coefficients $\{b_{l,\text{leg}}\}_{l=0}^{\infty}$, where $b_0 = 2/N$, is alternating and decreasing in magnitude. Furthermore, the ratios $\beta_l := |b_l/b_{l-1}|$, $l = 1, 2, \dots$ form a decreasing sequence with limiting value $\tilde{\beta} := \lim_{l \rightarrow \infty} \beta_l \geq b_0\beta_1$.¹² Because $\partial Q_p/\partial \phi > 0$ in the range $\phi \in (0, \pi)$, Q_p is bounded from above by $[2(1 - \beta_1)^{-1} - 1]/N$ and from below by $1/N - 2\beta_1/(N + 2\beta_1)$. For finite values of ε_J the ratios β_l and $\tilde{\beta}/\beta_1$ are monotonously increasing functions of ε_J with limiting value of unity, because $Q_p(\phi = 0, \varepsilon_J \rightarrow \infty) \searrow 0$, which enforces a stricter limit for the ratio $\tilde{\beta}/\beta_1$. This behaviour is explained by increasing long-range (high- l') correlations in Eq. (12) or actually in the state itself. On the other hand, the ratios are monotonously decreasing functions of N for fixed ε_J as correlations are weakened in longer arrays. All of the above-mentioned features are clearly shown in Table II, where β_l for the homogeneous case are depicted as function of N at $\varepsilon_J = 0.4$ and as function of ε_J for $N = 4$.¹³ These properties of the pumped charge are rather robust against relatively small systematical errors.

For small values of $\varepsilon_J \ll 1$ one has $\beta_{l>1} \approx b_0\beta_1$ and $\beta_1 \sim (N\varepsilon_J/2)^{N-2}[N(N-1)/2(N-2)!]$. The effects due to inhomogeneity, that is the relative sizes of $b_{1,\text{leg}}$, can be estimated by fixing the leg index r in the inhomogeneity prediction, Eq. (37) of Ref. 6. The corresponding results work quite well, only slightly overestimating the ratio between largest and smallest coefficients, even for large inhomogeneities X_{inh} . The total trans-

ferred charge $Q_p = \sum_{\text{leg}} Q_{p,\text{leg}}$ for inhomogeneous arrays does not have to satisfy $\beta_l > \beta_{l+1}$, although always $-b_{l+1}/b_l < 1$. This symmetry is also broken by non-ideal gating sequences, even for single legs.

In order to conclude, the properties of the tunnelling-charging Hamiltonian of a Cooper pair pump have been examined using an efficient block-diagonalisation scheme and a compatible Fourier expansion of the eigenstates. Explicit enforcement of the model symmetries produces strong systematics of the pumped charge, even if the structure of the Fourier coefficients $\{b_{l,\text{leg}}\}_{l=0}^{\infty}$ was not exhaustively proven. These properties are possibly related to the dynamical algebra of single and coupled Josephson junctions described in Ref. 14, which offers a complementary view of the present problem in case of a superconducting loop.

ACKNOWLEDGMENTS

This work has been supported by the Academy of Finland under the Finnish Centre of Excellence Programme 2000-2005 (Project No. 44875, Nuclear and Condensed Matter Programme at JYFL). The author thanks Mr. J.J. Toppari for insightful comments.

- ¹ H. Pothier, P. Lafarge, C. Urbina, D. Esteve and M.H. Devoret, *Europhys. Lett* **17**, 249 (1992).
- ² M. W. Keller, J. M. Martinis, N. M. Zimmerman and A.H. Steinbach, *Appl. Phys. Lett.* **69**, 1804 (1996); M.W. Keller, J.M. Martinis and R.L. Kautz, *Phys. Rev. Lett.* **80**, 4530 (1998); M. W. Keller, A. L. Eichenberger, J. M. Martinis and N. M. Zimmerman, *Science* **285**, 1716 (1999).
- ³ D. V. Averin, *Solid State Commun.* **105**, 659 (1998).
- ⁴ A. Shnirman and G. Schön, *Phys. Rev. B* **57**, 15400 (1998).
- ⁵ J. P. Pekola, J. J. Toppari, M. Aunola, M. T. Savolainen and D. V. Averin, *Phys. Rev. B* **60**, R9931 (1999).
- ⁶ M. Aunola, J. J. Toppari and J. P. Pekola, *Phys. Rev. B* **62**, 1296 (2000).
- ⁷ J. P. Pekola and J. J. Toppari, submitted to *Phys. Rev. Lett.* (2000).
- ⁸ G.-L. Ingold and Yu. V. Nazarov, in *Single Charge Tunnelling, Coulomb Blockade Phenomena in Nanostructures*, eds. H. Grabert and M.L. Devoret, (Plenum Press, New York, 1992).
- ⁹ W. Rudin, *Functional Analysis*, (McGraw Hill, 1973).
- ¹⁰ M. V. Berry, *Proc. R. Soc. Lond. A* **392**, 45 (1984).
- ¹¹ G. Falci, R. Fazio, G.M. Palma, J. Siewert, and V. Vedral, *Nature* **407**, 355 (2000).
- ¹² To some degree already in matrix elements $\langle m|\hat{M}|dm\rangle$.
- ¹³ The full data set and some tools for reproducing the data is available at <http://www.jyu.fi/~mimaau/Cooper/>
- ¹⁴ E. Celeghini, L. Faoro, and M. Rasetti, *Phys. Rev. B* **62**, 3054 (2000).